

Letter to the Editor

Mapping of the Loci for Mental Retardation Syndromes in the Distal Xq

To the Editor:

We constructed a deletion map of the distal Xq, using previously reported male interstitial deletions within Xq27–Xqter. The map consists of 12 large interstitial deletions, ranging from 100–9,000 kb. All but one fall within Xq27→proximal Xq28, forming a “deletion contig” between DXS51 (Xq26.3) and DXS455 (proximal 1/3 of Xq28). The region contains only a few genes, most of which are implicated in mental retardation (MR). Some of the deletions span more than one MR locus. The phenotype of affected males is typically limited to mental impairment unless the F9, IDS, or F8 gene is included in the deletion.

In the distal ⅓ of Xq28 only one large deletion of about 300 kb has been reported, and those remaining are small, intragenic deletions of <1 kb. This indicates that most deletions in this region are male-lethal.

The map allows us to narrow the location of several loci for syndromic MR [Neri et al., 1994], and facilitates mapping of new genes in the distal Xq.

Recent work at the junction of Xq27.3 and Xq28 demonstrates that this small region contains several loci implicated in mental retardation (MR): fragile-site mental retardation 1 (FMR1 and FRAXA) [Oostra and Verkerk, 1992]; fragile-site mental retardation 2 (FMR2 and FRAXE) [Knight et al., 1993; Gecz et al., 1996; Gu et al., 1996]; a locus around DXS296 [Gedeon et al., 1995] which may be the same as FMR2 [Gu et al., 1996]; iduronate 2-sulfatase (IDS) [Wilson et al., 1990, 1991]; and fragile-site F (FRAXF) [Hirst et al., 1993; Ritchie et al., 1994]. Thus, an abundance of MR loci is being found in the vast and otherwise void area of 15 Mb comprising Xq27→proximal Xq28, where only two other genes (cerebellar degeneration-

related autoantigen 1 (CDR1) [Hirst et al., 1991], and myotubular myopathy 1 (MTM1) [Dahl et al., 1995] have been assigned. It remains to be seen whether this clustering of MR genes reflects local peculiarities of the DNA sequence, or an ascertainment bias caused by the specific search for MR genes in the vicinity of the FMR1 gene. It is noteworthy that in the entire region Xq27–Xqter, every deletion found in a male is associated with mental retardation. Furthermore, some phenotypes may result from the absence of more than one MR gene. It should be considered, therefore, that the region Xq27→proximal Xq28 seems to be devoid of vital genes, and thus even very large deletions may produce no specific phenotype. Consequently, it is conceivable that normal individuals may also carry and propagate sizable deletions in Xq27 → proximal Xq28.

The situation is strikingly different in the gene-rich distal part of Xq28, where only one male deletion appears to cover more than one gene [Kenwrick et al., 1992]. Keeping in mind that females with deletions of Xq28 are fertile [Skibsted et al., 1984; Tharapel et al., 1993], the absence of such deletions in males suggests that they are male-lethal.

The positions and extent of the male deletions reported within Xq27–q28 are presented in Figure 1. Figure 1D depicts possible localization of the loci for syndromic MR whose genes have not yet been cloned. The location of the locus MRX27 [Glass et al., 1991] has been recently adjusted by Gedeon et al. [1996]. Map positions of the locus MRX3 [Gedeon et al., 1991] and the locus postulated by Kondo et al. [1991] could not be altered in this study, as both these conditions manifest as MR only. However, the positions of the genes involved in several MR syndromes can be considerably narrowed, because once a syndrome-specific phenotype is absent in a deleted patient, the respective gene can be excluded from the area of the deletion. Using these criteria, the positions of ANOP1, BD, DKC, IP2, MRSD, OPD1, WSN, BFLS, and ADS can be narrowed down, as depicted on the map (Fig. 1D). The exclusion map presented will facilitate clinical interpretation of contiguous gene syndromes, as well as localization of new loci in the distal Xq.

Received for publication October 24, 1995; revision received December 18, 1995.

Address reprint requests to Malgorzata Schmidt, Department of Genetics and Human Variation, La Trobe University, Melbourne 3083, Australia.




Fig. 1. Male deletions in the distal Xq. **A:** Map of the X chromosome scaled in megabases. **B:** Deletions reported in males are represented by vertical lines, with name of first author and approximate size in kb. **C:** Order of X chromosome markers [Willard et al., 1994.] Bold horizontal symbols indicate position of known genes involved in mental retardation. The position of SOX3 [Stevanovic et al., 1993] has been narrowed (Schmidt, unpublished results) using an X chromosome with a duplication DXS152→Xqter [Schmidt et al., 1991]. **D:** Potential location of loci for the mental retardation syndromes listed below: ANOP1, anophthalmos (Graham syndrome) MIM 301590; ADS, Ataxia-dementia syndrome MIM 301840; BFLS, Börjeson-Forssman-Lehmann syndrome MIM 301900; BD, bullous dystrophy MIM 302000; DKC, dyskeratosis congenita MIM 305000; IP2, incontinentia pigmenti type II MIM 308310; MRSD, mental retardation, skeletal dysplasia (Christian syndrome) MIM 309620; OPD1, otopalatodigital syndrome MIM 311300; WSN, Waisman syndrome (Parkinsonism, Laxova syndrome) MIM 311510. References: Fuchs <1, Sarde et al., 1994; Gedeon 200, 100: Gedeon et al., 1995; Gedeon 2500, Gedeon et al., 1992; MRX3, Gedeon et al., 1991; MRX27, Glass et al., 1991; Gedeon et al., 1996; MTM1, Dahl et al., 1995; SOX3, Stevanovic et al., 1993; Tarleton ~ 5000, Albright et al., 1994; Wilson 1700, 800, Steen-Bondeson et al., 1992; Wohrle 250-Hirst 25; 0.6, Hirst et al., 1995 (deletions reported by de Graaff et al. [1995] are not included in Figure 1, as they result from somatic instability of the expanded FMR1 gene).

REFERENCES

- Albright SG, Lachiewicz AM, Tarleton JC, Rao KW, Schwartz CE, Richie R, Tennison MB, Aylsworth AS (1994): Fragile X phenotype in a patient with a large de novo deletion in Xq27-q28. *Am J Med Genet* 51:294-297.
- Beck M, Steglich C, Zabel B, Dahl N, Schwinger E, Hopwood JJ (1992): Deletion of the Hunter gene and both DXS466 and DXS304 in a patient with mucopolysaccharidosis type II. *Am J Med Genet* 44:100-103.
- Dahl N, Hu LJ, Chery M, Fardeau M, Gilgenkrantz S, Nivelon-Chevallier A, Sidaner-Noisette I, Mugneret F, Gouyon J-B, Gal A, Kioschis P, d'Urso M, Mandel J-L (1995): Myotubular myopathy in a girl with a deletion at Xq27-q28 and unbalanced X inactivation assigns the MTM1 gene to a 600-kb region. *Am J Hum Genet* 56:1108-1115.
- De Graaff E, Rouillard P, Willems P, Smits APT, Rousseau F, Oostra BA (1995): Hotspot for deletions in the CGG repeat region of FMR1 in fragile X patients. *Hum Mol Genet* 4:45-49.
- Fuchs S, Sarde CO, Wedemann H, Schwinger E, Mendel J-L, Gal A (1994): Missense mutations are frequent in the gene for X-chromosomal adrenoleukodystrophy (ALD). *Hum Mol Genet* 3:1903-1905.
- Gecz J, Gedeon AK, Sutherland GR, Mulley JC (1996): Identification of the gene FMR2, associated with the FRAXE mental retardation. *Nat Genet* 13:105-108.
- Gedeon AK, Kerr B, Mulley J, Turner G (1991): Localization of the MRX3 gene for nonspecific X-linked mental retardation. *J Med Genet* 28:372-377.
- Gedeon AK, Baker E, Robinson H, Partington MW, Gross B, Manca A, Korn B, Poustka A, Yu S, Sutherland GR, Mulley JC (1992): Fragile X syndrome without CCG amplification has an FMR1 deletion. *Nat Genet* 1:341-344.
- Gedeon AK, Glass IA, Connor JM, Mulley JC (1996): Localization of MRX27 to Xq24-q26 delineates the sixth discrete gene for nonspecific X-linked mental retardation. *Am J Med Genet* 64:121-124.
- Gedeon AK, Keinanen M, Ades LC, Kaariainen H, Gecz J, Baker E, Sutherland GR, Mulley JC (1995): Overlapping submicroscopic deletions in Xq28 in two unrelated boys with developmental disorders: Identification of a gene near FRAXE. *Am J Hum Genet* 56:907-914.
- Glass IA, White EM, Pope MJ, Pirrit LA, Cockburn F, Connor JM (1991): Linkage analysis in a large family with nonspecific X-linked mental retardation. *Am J Med Genet* 38:240-243.
- Gu Y, Shen Y, Gibbs RA, Nelson DL (1996): Identification of FMR2, a novel gene associated with the FRAXE CCG repeat and CpG island. *Nat Genet* 13:109-113.
- Gu Y, Lugenbeel KA, Vockley JG, Grody WW, Nelson DL (1994): A de novo deletion in FMR1 in a patient with developmental delay. *Hum Mol Genet* 3:1705-1706.
- Hirst MC, Barnicoat A, Flynn G, Wang Q, Daker M, Buckle VJ, Davies KE, Bobrow M (1993): The identification of a third fragile site, FRAXF in Xq27-28 distal to both FRAXA and FRAXE. *Hum Mol Genet* 2:197-200.
- Hirst MC, Bell MV, MacKinnon RN, Watson JE, Callen D, Sutherland G, Dahl N, Patterson MN, Schwartz C, Ledbetter D (1991): Mapping of a cerebellar degeneration-related protein and DXS304 around the fragile site. *Am J Med Genet* 38:354-356.
- Hirst M, Grewal P, Flannery A, Slatter R, Maher E, Barton D, Fryns J-P, Davies K (1995): Two new cases of FMR1 deletion associated with mental impairment. *Am J Hum Genet* 56:67-74.
- Kenwrick S, Levinson B, Taylor S, Shapiro A, Gitschier J (1992): Isolation and sequence of two genes associated with a CpG island 5' of the factor VIII gene. *Hum Mol Genet* 1:179-186.
- Knight SJL, Flannery AV, Hirst MC, Campbell L, Christodoulou Z, Phelps SR, Pointon J, Middleton-Price HR, Barnicoat A, Pembrey ME, Holland J, Oostra BA, Bobrow M, Davies KE (1993): Trinucleotide repeat amplification and hypermethylation of a CpG island in FRAXE/mental retardation. *Cell* 74:127-134.
- Kondo I, Tsukamoto K, Niikawa N, Okano K, Kanazawa I, Hupkes PE (1991): A new form of X-linked mental retardation linked to DXS 369. *Cytogenet Cell Genet* 58:2071.
- Meijer H, de Graaff E, Merckx ML, Jongbloed JE, de Die-Smulders EM, Englen JJM, Fryns J-P, Curfs PMG, Oostra BA (1994): A deletion of 1.6 kb proximal to the CGG repeat of the FMR1 gene causes the clinical phenotype of the fragile X syndrome. *Hum Mol Genet* 3:615-620.
- Neri G, Chiurazzi P, Arena JF, Lubs HA (1994): XLMR genes: Update 1994. *Am J Med Genet* 51:542-549.
- Oostra B, Verkerk AJMH (1992): Review: The fragile X syndrome: Isolation of the FMR-1 gene and characterization of the fragile X mutation. *Chromosoma* 101:381-387.
- Palmieri G, Capra V, Romano G, D'Urso M, Johnson S, Schlessinger D, Morris P, Hopwood J, Di Natale P, Gatti R, Ballabio A (1992): The iduronate sulfatase gene: Isolation of a 1.2-Mb YAC contig spanning the entire gene and identification of heterogenous deletions in patients with Hunter syndrome. *Genomics* 12:52-57.
- Quan F, Zonana J, Gunter K, Peterson KL, Magenis RE, Popovich BW (1995): An atypical case of fragile X syndrome caused by a deletion that includes the FMR1 gene. *Am J Hum Genet* 56:1042-1051.
- Ritchie RJ, Knight SJL, Hirst MC, Grewal PK, Bobrow M, Cross GS, Davies KE (1994): The cloning of FRAXF: Trinucleotide repeat expansion and methylation at a third fragile site in distal Xqter. *Hum Mol Genet* 3:2115-121.
- Rosenthal A, Jouet M, Kenwrick S (1992): Aberrant splicing of neural cell adhesion molecule L1 mRNA in a family with X-linked hydrocephalus. *Nat Genet* 2:107-112.
- Rousseau F, Vincent A, Rivella S, Heitz D, Triboli C, Maestrini E, Warren ST, Suthers GK, Goodfellow P, Mandel J-L, Toniolo D, Oberle I (1991): Four chromosomal breakpoints and four new probes mark out a 10-cM region encompassing the fragile X locus (FRAXA). *Am J Hum Genet* 48:108-116.
- Sarde C-O, Mosser J, Kioschis P, Kretz C, Vicaire S, Aubourg P, Poustka A, Mandel J-L (1994): Genomic organization of the adrenoleukodystrophy gene. *Genomics* 22:13-20.
- Schmidt M, Du Sart D, Kalitsis P, Leversha M, Dale S, Sheffield L, Toniolo D (1991): Duplications of the X chromosome in males: Evidence that most parts of the X chromosome can be active in two copies. *Hum Genet* 86:519-521.
- Skibsted L, Westh H, Niebuhr E (1984): X long-arm deletions. A review of non-mosaic cases studied with banding techniques. *Hum Genet* 67:1-5.
- Steen-Bondeson M-L, Dahl N, Tonnesen T, Kleijer WJ, Seidlitz G, Gustavson K-H, Wilson PJ, Morris CP, Hopwood JJ, Pettersson U (1992): Molecular analysis of patients with Hunter syndrome: Implication of a region prone to structural alterations within the IDS gene. *Hum Mol Genet* 1:195-198.
- Stevanovic M, Lovell-Badge R, Collignon J, Goodfellow PN (1993): SOX3 is an X-linked gene related to SRY. *Hum Mol Genet* 12:2013-2018.
- Tarleton JC, Richie R, Schwartz C, Rao K, Aylsworth AS, Lachiewicz A (1993): An extensive de novo deletion removing FMR1 in a patient with mental retardation and the fragile X syndrome phenotype. *Hum Mol Genet* 2:1973-1974.
- Tharapel AT, Anderson KP, Simpson JL, Martens PR, Wilroy RS, Llerena JC, Schwartz CE (1993): Deletion (X)(q26.1→q28) in a proband and her mother: Molecular characterization and phenotypic-karyotypic deductions. *Am J Hum Genet* 52:463-471.
- Trottier Y, Imbert G, Poustka A, Fryns J-P, Mandel J-L (1994): Male with typical fragile X phenotype is deleted for part of the FMR1 gene and for about 100 kb of upstream region. *Am J Med Genet* 51:454-457.
- Willard HF, Cremers F, Mandel J-L, Monaco AP, Nelson DL, Schlessinger D (1994): Report of the Fifth International Workshop on Human X Chromosome Mapping 1994. *Cytogenet Cell Genet* 67:296-328.
- Wilson PJ, Morris CP, Anson DS, Occhiodoro T, Bielicki J, Clements PR, Hopwood JJ (1990): Hunter syndrome: Isolation of an iduronate-2-sulfatase cDNA clone and analysis of patient DNA. *Proc Natl Acad Sci USA* 87:8531-8535.
- Wilson PJ, Suthers GK, Callen DF, Baker E, Nelson PV, Cooper A, Wraith JE, Sutherland GR, Morris CP, Hopwood JJ (1991): Frequent deletions at Xq28 indicate genetic heterogeneity in Hunter syndrome. *Hum Genet* 86:505-508.
- Wohrle D, Kotzot D, Hirst MC, Manca A, Korn B, Schmidt A, Barbi G, Rott H-D, Poustka A, Davies KE, Steinbach P (1992): A microdeletion of less than 250 kb, including the proximal part of the FMR-1 gene and the fragile X site, in a male with the clinical phenotype of fragile-X syndrome. *Am J Hum Genet* 51:299-306.

Malgorzata Schmidt

Department of Genetics and Human Variation
La Trobe University
Melbourne, Australia